Causes of Extreme Dry Conditions over Western U.S. during Early 2013

Hailan Wang¹² and Siegfried Schubert¹ NASA/GMAO¹; SSAI²

October 21, 2013





Extreme Dry Anomalies over Western Coastal U.S.


NCDC (wrt 1895-present)

January-March 2013 Statewide Ranks National Climatic Data Center/NESDIS/NOAA

20 49 19 3 58 71 44 32 78 106 75 87 51 31 55 Precipitation 1 = Driest 119 = Wettest Record Much Below Near Above Much Record Driest Normal Above Wettest Normal Normal


Normal

NOAA PREC (wrt 1980-2010)




Precip over Western U.S.

- Mainly over Pacific northwest, and comes from extratropical cyclones from the north Pacific under the strong wintertime jet stream
- Precip over California: southwesterlies from north of Hawaii
- Influencing factors
 - ENSO
 - MJO
 - Atmospheric internal variability
 - Variations on decadal and longer time scales
 - · PDO
 - · Long-term trend




Data


- Obs precip
 - NOAA PREC (Jan1948-Apr2013)
 - GPCP (Jan1979-Dec2011)
- MERRA reanalysis
 - Jan1979-present
- NASA GEOS-5 AGCM Simulations
 - 1 degree
 - AMIP
 - 12 members; Jan1871-present
 - Idealized AGCM runs
 - Clim SST (1901-2004)
 - Anom SST: ColdPac+WarmAtl+WarmTrend
 - 50 years long, with a repeating seasonal cycle
- Anomalies: wrt 1980-2010 mean



ColdPac + WarmAtl + WarmTrend





MERRA: Daily Evolution



Considerably less north Pacific storms reached U.S. west coast Considerably weaker zonal wind over NE Pacific

Atmospheric Circulation: 250mb; JF2013








Similar to other dry winters for California, persistent ridge (40°N) and weaker zonal wind (30°N) over NE Pacific prevented north Pacific storms from reaching U.S. west coast.

SSTA











Cold ENSO like pattern, but with weak SST indices



0.5

1.5


Precip: AMIP EnsMean vs. MERRA



SSTA in Pacific contributed to the dry anomalies over SW US

Stationary Wave Modeling Diagnosis

Eddy u250; JF2013; MERRA





The weakened zonal wind over NE Pacific was primarily maintained by changes in transient vorticity flux convergences (TranVor), and secondly by heating (Heat) anomalies.

Climate Variations on Decadal and Longer Time Scale

Through Changing Basic State

Clim vs Anom (ColdPac+WarmAtl+WarmTrend)






PcAwLTw dominated by Cold Pacific

The effect of PcAwLTw through changing mean basic state is investigated by examining the changes in optimal forcing pattern for the zonal wind over NE Pacific, using a stationary wave model

Climate Variations on Decadal and Longer Time Scale Optimal Forcing Pattern for U250 over NE Pacific



The changes to the basic state due to PcAwLTw allows forcing over Pacific to exert stronger effects over NE Pacific and western U.S.

Summary

Immediate Cause

 Less North Pacific storms reached west coast of U.S., due to considerably weaker zonal wind over NE Pacific

Underlying Causes

- SSTA: cold ENSO-like SSTA contributed to dry anomalies over SW U.S.
- Atmospheric internal variability: dominant role of transient vorticity flux convergences for the weakened zonal wind over NE Pacific
- Climate variations on decadal and longer time scales, especially the ongoing negative PDO: the changes to the basic state allows forcing over Pacific to exert stronger effects over NE Pacific and western U.S.